

Hermite-Hadamard type inequalities for product of GA-convex functions via Hadamard fractional integrals

İmdat İşcan and Mehmet Kunt

Abstract. In this paper, some Hermite-Hadamard type inequalities for products of two GA-convex functions via Hadamard fractional integrals are established. Our results about GA-convex functions are analogous generalizations for some other results proved by Pachpette for convex functions.

Mathematics Subject Classification (2010): 26A51, 26A33, 26D10.

Keywords: Hermite-Hadamard inequality, GA-convex functions, Hadamard fractional integral.

1. Introduction

In recent years, very large number of studies of error estimations have been done for Hermite-Hadamard type inequalities. It is known that Hermite-hadamard integral inequality was built on a convex function. In time, Hermite-Hadamard inequality is developed other kinds of convex functions. For some results which generalize, improve, and extend the Hermite-Hadamard inequality see [1, 7, 10, 18, 20] and references therein.

Hermite-Hadamard type inequalities for products of two convex functions are interesting problem and firstly developed by Pachpette in [16]. In [17], Pachpette also established Hermite-hadamard type inequalities involving two log-convex functions. In [11], Kırmacı et. al. proved several Hermite-Hadamard type inequalities for products of two convex and s -convex functions. In [19], Sarıkaya et. al. proved some Hermite-Hadamard type inequalities for products of two h -convex functions. In [2], Bakula et. al. established Hermite-Hadamard type inequalities for products of two m -convex and (α, m) -convex functions. In [4, 6], Chen and Wu obtained some Hermite-Hadamard type inequalities for products of two convex and harmonically s -convex functions. In [21], Yin and Qi established some Hermite-Hadamard type inequalities for products

of two convex functions. In [5], Chen obtained some new Hermite-Hadamard type inequalities for products of two convex functions via Riemann-Liouville fractional integrals and in [3] he extended this problem to m -convex and (α, m) -convex functions.

In this work, we establish Hermite-Hadamard type inequalities for products of two GA-convex functions via Hadamard fractional integrals. Our results are analogous generalization for some results in [16].

2. Preliminaries

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \tag{2.1}$$

is well known in the literature as Hermite-Hadamard's inequality [8].

In [16], Pachpette established following two Hermite-Hadamard type inequalities for products of convex functions as follows:

Theorem 2.1. *Let f and g be real-valued, non-negative and convex functions on $[a, b]$. Then*

$$\frac{1}{b-a} \int_a^b f(x)g(x)dx \leq \frac{1}{3}M(a,b) + \frac{1}{6}N(a,b) \tag{2.2}$$

and

$$2f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)g(x)dx + \frac{1}{6}M(a,b) + \frac{1}{3}N(a,b) \tag{2.3}$$

where $M(a,b) = f(a)g(a) + f(b)g(b)$ and $N(a,b) = f(a)g(b) + f(b)g(a)$.

Definition 2.2. [14, 15]. A function $f : I \subseteq (0, \infty) \rightarrow \mathbb{R}$ is said to be GA-convex (geometric-arithmetically convex) if

$$f(x^t y^{1-t}) \leq tf(x) + (1-t)f(y)$$

for all $x, y \in I$ and $t \in [0, 1]$.

We will now give definitions of the right-hand side and left-hand side Hadamard fractional integrals which are used throughout this paper.

Definition 2.3. [12]. Let $f \in L[a, b]$. The right-hand side and left-hand side Hadamard fractional integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ of order $\alpha > 0$ with $b > a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x \left(\ln \frac{x}{t}\right)^{\alpha-1} f(t) \frac{dt}{t}, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b \left(\ln \frac{t}{x}\right)^{\alpha-1} f(t) \frac{dt}{t}, \quad x < b$$

respectively, where $\Gamma(\alpha)$ is the Gamma function defined by

$$\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt.$$

In [9], İşcan represented Hermite-Hadamard's inequalities for GA-convex functions in fractional integral forms as follows.

Theorem 2.4. *Let $f : I \subseteq (0, \infty) \rightarrow \mathbb{R}$ be a function such that $f \in L[a, b]$ where $a, b \in I$ with $a < b$. If f is a GA-convex function on $[a, b]$, then the following inequalities for fractional integrals hold:*

$$f(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2(\ln \frac{b}{a})^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] \leq \frac{f(a) + f(b)}{2} \tag{2.4}$$

with $\alpha > 0$.

In [13], Kunt and İşcan established new Hermite-Hadamard type inequality for GA-convex function in fractional integral forms as follows:

Theorem 2.5. *Let $f : [a, b] \subseteq (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function with $a < b$ and $f \in L[a, b]$, then the following inequalities for fractional integrals hold:*

$$f(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2^{1-\alpha}(\ln \frac{b}{a})^\alpha} [J_{\sqrt{ab}-}^\alpha f(a) + J_{\sqrt{ab}+}^\alpha f(b)] \leq \frac{f(a) + f(b)}{2}. \tag{2.5}$$

3. General results

Theorem 3.1. *Let f and $g : [a, b] \subseteq (0, \infty) \rightarrow \mathbb{R}$ be non-negative and GA-convex functions with $a < b$ and $f \in L[a, b]$, then the following inequality for fractional integrals hold:*

$$\begin{aligned} & \frac{\Gamma(\alpha + 1)}{2(\ln \frac{b}{a})^\alpha} [J_{a+}^\alpha f(b)g(b) + J_{b-}^\alpha f(a)g(a)] \\ & \leq \left(\frac{\alpha}{\alpha + 2} - \frac{\alpha}{\alpha + 1} + \frac{1}{2} \right) M(a, b) + \frac{\alpha}{(\alpha + 2)(\alpha + 1)} N(a, b) \end{aligned} \tag{3.1}$$

where $\alpha > 0$, $M(a, b) = f(a)g(a) + f(b)g(b)$ and $N(a, b) = f(a)g(b) + f(b)g(a)$.

Proof. Since f and g are non-negative and GA-convex functions on $[a, b]$, we have for all $t \in [0, 1]$

$$f(a^t b^{1-t}) \leq tf(a) + (1-t)f(b), \tag{3.2}$$

and

$$g(a^t b^{1-t}) \leq tg(a) + (1-t)g(b). \tag{3.3}$$

From products of (3.2) and (3.3), we have

$$\begin{aligned} f(a^t b^{1-t})g(a^t b^{1-t}) & \leq t^2 f(a)g(a) + (1-t)^2 f(b)g(b) \\ & \quad + t(1-t)[f(a)g(b) + f(b)g(a)]. \end{aligned} \tag{3.4}$$

Similarly (3.4), we have

$$f(a^{1-t}b^t)g(a^{1-t}b^t) \leq (1-t)^2 f(a)g(a) + t^2 f(b)g(b) + t(1-t)[f(a)g(b) + f(b)g(a)]. \tag{3.5}$$

The sum of (3.4) and (3.5), we have

$$f(a^t b^{1-t})g(a^t b^{1-t}) + f(a^{1-t} b^t)g(a^{1-t} b^t) \leq (2t^2 - 2t + 1) M(a, b) + 2t(1-t) N(a, b) \tag{3.6}$$

Multiplying both sides of (3.6) by $t^{\alpha-1} \frac{\alpha}{2}$, then integrating the obtained inequality with respect to t over $[0, 1]$, we have

$$\begin{aligned} & \frac{\alpha}{2} \left[\int_0^1 t^{\alpha-1} f(a^t b^{1-t})g(a^t b^{1-t})dt + \int_0^1 t^{\alpha-1} f(a^{1-t} b^t)g(a^{1-t} b^t)dt \right] \\ &= \frac{\alpha}{2} \left[\int_a^b \left(\frac{\ln \frac{b}{u}}{\ln \frac{b}{a}} \right)^{\alpha-1} f(u)g(u) \frac{du}{u \ln \frac{b}{a}} + \int_a^b \left(\frac{\ln \frac{v}{a}}{\ln \frac{b}{a}} \right)^{\alpha-1} f(v)g(v) \frac{du}{v \ln \frac{b}{a}} \right] \\ &= \frac{\alpha}{2 \left(\ln \frac{b}{a} \right)^\alpha} \left[\int_a^b \left(\ln \frac{b}{u} \right)^{\alpha-1} f(u)g(u) \frac{du}{u} + \int_a^b \left(\ln \frac{v}{a} \right)^{\alpha-1} f(v)g(v) \frac{du}{v} \right] \\ &= \frac{\Gamma(\alpha + 1)}{2 \left(\ln \frac{b}{a} \right)^\alpha} [J_{a+}^\alpha f(b)g(b) + J_{b-}^\alpha f(a)g(a)] \\ &\leq \frac{\alpha}{2} \left[M(a, b) \int_0^1 t^{\alpha-1} (2t^2 - 2t + 1) dt + N(a, b) \int_0^1 t^{\alpha-1} 2t(1-t) dt \right] \\ &= \left(\frac{\alpha}{\alpha + 2} - \frac{\alpha}{\alpha + 1} + \frac{1}{2} \right) M(a, b) + \frac{\alpha}{(\alpha + 2)(\alpha + 1)} N(a, b) \end{aligned}$$

and this completes the proof. □

Remark 3.2. Theorem 3.1 is an analogous generalization of (2.2) for GA-convex functions.

Corollary 3.3. In Theorem 3.1, if we take $g : [a, b] \rightarrow \mathbb{R}$ as $g(x) = 1$ for all $x \in [a, b]$, then we have

$$\frac{\Gamma(\alpha + 1)}{2 \left(\ln \frac{b}{a} \right)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] \leq \frac{f(a) + f(b)}{2}$$

which is the right hand side of (2.4).

Corollary 3.4. In Theorem 3.1, if we take $\alpha = 1$, then we have

$$\frac{1}{\ln b - \ln a} \int_a^b f(x)g(x) \frac{dx}{x} \leq \frac{1}{3}M(a, b) + \frac{1}{6}N(a, b)$$

for GA-convex functions.

Theorem 3.5. *Let f and $g : [a, b] \subseteq (0, \infty) \rightarrow \mathbb{R}$ be non-negative and GA-convex functions with $a < b$ and $f \in L[a, b]$, then the following inequality for fractional integrals hold:*

$$2f(\sqrt{ab})g(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2(\ln \frac{b}{a})^\alpha} [J_{a+}^\alpha f(b)g(b) + J_{b-}^\alpha f(a)g(a)] + \frac{\alpha}{(\alpha + 2)(\alpha + 1)}M(a, b) + \left(\frac{\alpha}{\alpha + 2} - \frac{\alpha}{\alpha + 1} + \frac{1}{2}\right)N(a, b) \tag{3.7}$$

where $\alpha > 0$, $M(a, b) = f(a)g(a) + f(b)g(b)$ and $N(a, b) = f(a)g(b) + f(b)g(a)$.

Proof. It is clear for all $t \in [0, 1]$

$$\sqrt{ab} = \sqrt{a^t b^{1-t} \cdot a^{1-t} b^t} = \sqrt{a^t b^{1-t}} \sqrt{a^{1-t} b^t}.$$

Since f and g are non-negative and GA-convex functions on $[a, b]$, we have for all $t \in [0, 1]$

$$\begin{aligned} f(\sqrt{ab})g(\sqrt{ab}) &= f(\sqrt{a^t b^{1-t}} \sqrt{a^{1-t} b^t})g(\sqrt{a^t b^{1-t}} \sqrt{a^{1-t} b^t}) \\ &\leq \frac{1}{4} [f(a^t b^{1-t}) + f(a^{1-t} b^t)] [g(a^t b^{1-t}) + g(a^{1-t} b^t)] \\ &= \frac{1}{4} [f(a^t b^{1-t})g(a^t b^{1-t}) + f(a^{1-t} b^t)g(a^{1-t} b^t)] \\ &\quad + \frac{1}{4} [f(a^t b^{1-t})g(a^{1-t} b^t) + f(a^{1-t} b^t)g(a^t b^{1-t})] \\ &\leq \frac{1}{4} [f(a^t b^{1-t})g(a^t b^{1-t}) + f(a^{1-t} b^t)g(a^{1-t} b^t)] \\ &\quad + \frac{1}{4} [tf(a) + (1-t)f(b)] [(1-t)g(a) + tg(b)] \\ &\quad + \frac{1}{4} [(1-t)f(a) + tf(b)] [tg(a) + (1-t)g(b)] \\ &= \frac{1}{4} [f(a^t b^{1-t})g(a^t b^{1-t}) + f(a^{1-t} b^t)g(a^{1-t} b^t)] \\ &\quad + \frac{1}{4} \{2t(1-t)[f(a)g(a) + f(b)g(b)] \\ &\quad + (2t^2 - 2t + 1)[f(a)g(b) + f(b)g(a)]\} \end{aligned} \tag{3.8}$$

Multiplying both sides of (3.8) by $2\alpha t^{\alpha-1}$, then integrating the obtained inequality with respect to t over $[0, 1]$, we have

$$2f(\sqrt{ab})g(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2(\ln \frac{b}{a})^\alpha} [J_{a+}^\alpha f(b)g(b) + J_{b-}^\alpha f(a)g(a)] + \frac{\alpha}{(\alpha + 2)(\alpha + 1)}M(a, b) + \left(\frac{\alpha}{\alpha + 2} - \frac{\alpha}{\alpha + 1} + \frac{1}{2}\right)N(a, b)$$

and this completes the proof. □

Remark 3.6. Theorem 3.5 is an analogous generalization of (2.3) for GA-convex functions.

Corollary 3.7. *In Theorem 3.5, if we take $g : [a, b] \rightarrow \mathbb{R}$ as $g(x) = 1$ for all $x \in [a, b]$, then we have*

$$2f(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2\left(\ln \frac{b}{a}\right)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] + \frac{f(a) + f(b)}{2}.$$

Corollary 3.8. *In Theorem 3.5, if we take $\alpha = 1$, then we have*

$$2f(\sqrt{ab})g(\sqrt{ab}) \leq \frac{1}{\ln b - \ln a} \int_a^b f(x)g(x) \frac{dx}{x} + \frac{1}{6}M(a, b) + \frac{1}{3}N(a, b)$$

for GA-convex functions.

Theorem 3.9. *Let f and $g : [a, b] \subseteq (0, \infty) \rightarrow \mathbb{R}$ be non-negative and GA-convex functions with $a < b$ and $f \in L[a, b]$, then the following inequality for fractional integrals hold:*

$$\begin{aligned} & \frac{\Gamma(\alpha + 1)}{2^{1-\alpha} \left(\ln \frac{b}{a}\right)^\alpha} \left[J_{\sqrt{ab}^-}^\alpha f(a)g(a) + J_{\sqrt{ab}^+}^\alpha f(b)g(b) \right] \\ & \leq \left(\frac{\alpha}{4(\alpha + 2)} - \frac{\alpha}{2(\alpha + 1)} + \frac{1}{2} \right) M(a, b) + \frac{\alpha^2 + 3\alpha}{4(\alpha + 2)(\alpha + 1)} N(a, b) \end{aligned} \quad (3.9)$$

where $\alpha > 0$, $M(a, b) = f(a)g(a) + f(b)g(b)$ and $N(a, b) = f(a)g(b) + f(b)g(a)$.

Proof. Since f and g are non-negative and GA-convex functions on $[a, b]$, multiplying both sides of (3.6) by $t^{\alpha-1} \frac{\alpha}{2^{1-\alpha}}$, then integrating the obtained inequality with respect to t over $[0, \frac{1}{2}]$, we have

$$\begin{aligned} & \frac{\alpha}{2^{1-\alpha}} \left[\int_0^{\frac{1}{2}} t^{\alpha-1} f(atb^{1-t})g(atb^{1-t})dt + \int_0^{\frac{1}{2}} t^{\alpha-1} f(a^{1-t}b^t)g(a^{1-t}b^t)dt \right] \\ & = \frac{\alpha}{2^{1-\alpha}} \left[\int_{\sqrt{ab}}^b \left(\frac{\ln \frac{b}{u}}{\ln \frac{b}{a}} \right)^{\alpha-1} f(u)g(u) \frac{du}{u \ln \frac{b}{a}} + \int_a^{\sqrt{ab}} \left(\frac{\ln \frac{v}{a}}{\ln \frac{b}{a}} \right)^{\alpha-1} f(v)g(v) \frac{du}{v \ln \frac{b}{a}} \right] \\ & = \frac{\alpha}{2^{1-\alpha} \left(\ln \frac{b}{a}\right)^\alpha} \left[\int_{\sqrt{ab}}^b \left(\ln \frac{b}{u} \right)^{\alpha-1} f(u)g(u) \frac{du}{u} + \int_a^{\sqrt{ab}} \left(\ln \frac{v}{a} \right)^{\alpha-1} f(v)g(v) \frac{du}{v} \right] \\ & = \frac{\Gamma(\alpha + 1)}{2^{1-\alpha} \left(\ln \frac{b}{a}\right)^\alpha} \left[J_{\sqrt{ab}^+}^\alpha f(b)g(b) + J_{\sqrt{ab}^-}^\alpha f(a)g(a) \right] \\ & \leq \frac{\alpha}{2^{1-\alpha}} \left[M(a, b) \int_0^{\frac{1}{2}} t^{\alpha-1} (2t^2 - 2t + 1) dt + N(a, b) \int_0^{\frac{1}{2}} t^{\alpha-1} 2t(1-t) dt \right] \\ & = \left(\frac{\alpha}{4(\alpha + 2)} - \frac{\alpha}{2(\alpha + 1)} + \frac{1}{2} \right) M(a, b) + \frac{\alpha^2 + 3\alpha}{4(\alpha + 2)(\alpha + 1)} N(a, b) \end{aligned}$$

and this completes the proof. □

Remark 3.10. Theorem 3.9 is an other analogous generalization of (2.2) for GA-convex functions.

Corollary 3.11. *In Theorem 3.9, if we take $g : [a, b] \rightarrow \mathbb{R}$ as $g(x) = 1$ for all $x \in [a, b]$, then we have*

$$\frac{\Gamma(\alpha + 1)}{2^{1-\alpha} \left(\ln \frac{b}{a}\right)^\alpha} \left[J_{\sqrt{ab}-}^\alpha f(a) + J_{\sqrt{ab}+}^\alpha f(b) \right] \leq \frac{f(a) + f(b)}{2}$$

which is the right hand side of (2.5).

Corollary 3.12. *In Theorem 3.9, if we take $\alpha = 1$, then we have*

$$\frac{1}{\ln b - \ln a} \int_a^b f(x) g(x) \frac{dx}{x} \leq \frac{1}{3} M(a, b) + \frac{1}{6} N(a, b)$$

for GA-convex functions.

Theorem 3.13. *Let f and $g : [a, b] \subseteq (0, \infty) \rightarrow \mathbb{R}$ be non-negative and GA-convex functions with $a < b$ and $f \in L[a, b]$, then the following inequality for fractional integrals hold:*

$$2f(\sqrt{ab}) g(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2^{1-\alpha} \left(\ln \frac{b}{a}\right)^\alpha} \left[J_{\sqrt{ab}-}^\alpha f(a) g(a) + J_{\sqrt{ab}+}^\alpha f(b) g(b) \right] + \frac{\alpha^2 + 3\alpha}{4(\alpha + 2)(\alpha + 1)} M(a, b) + \left(\frac{\alpha}{4(\alpha + 2)} - \frac{\alpha}{2(\alpha + 1)} + \frac{1}{2} \right) N(a, b) \quad (3.10)$$

where $\alpha > 0$, $M(a, b) = f(a) g(a) + f(b) g(b)$ and $N(a, b) = f(a) g(b) + f(b) g(a)$.

Proof. Multiplying both sides of (3.8) by $2^{1+\alpha} \alpha t^{\alpha-1}$, then integrating the obtained inequality with respect to t over $[0, \frac{1}{2}]$, we have desired result. \square

Remark 3.14. Theorem 3.13 is an other analogous generalization of (2.3) for GA-convex functions.

Corollary 3.15. *In Theorem 3.13, if we take $g : [a, b] \rightarrow \mathbb{R}$ as $g(x) = 1$ for all $x \in [a, b]$, then we have*

$$2f(\sqrt{ab}) \leq \frac{\Gamma(\alpha + 1)}{2^{1-\alpha} \left(\ln \frac{b}{a}\right)^\alpha} \left[J_{\sqrt{ab}-}^\alpha f(a) + J_{\sqrt{ab}+}^\alpha f(b) \right] + \frac{f(a) + f(b)}{2}.$$

Corollary 3.16. *In Theorem 3.13, if we take $\alpha = 1$, then we have*

$$2f(\sqrt{ab}) g(\sqrt{ab}) \leq \frac{1}{\ln b - \ln a} \int_a^b f(x) g(x) \frac{dx}{x} + \frac{1}{6} M(a, b) + \frac{1}{3} N(a, b)$$

for GA-convex functions.

References

- [1] Bai, S.P., Wang, S.H., Qi, F., *Some Hermite-Hadamard type inequalities for n-time differentiable (α, m) -convex functions*, J. Inequal. Appl., **267**(2012), 11 pages.
- [2] Bakula, M.K., Özdemir, M.E., Pečarić, J., *Hadamard type inequalities form m -convex and (α, m) -convex functions*, Journal of Inequalities in Pure and Applied Mathematics, **9**(2008), no. 4, article 96.

- [3] Chen, F., *On Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals via two kinds of convexity*, Chinese Journal of Mathematics, Volume 2014, Article ID:173293, 2014.
- [4] Chen, F., *A note on Hermite-Hadamard inequalities for products of convex functions*, Journal of Applied Mathematics, Volume 2013, Article ID:935020, 2013.
- [5] Chen, F., *A note on Hermite-Hadamard inequalities for products of convex functions via Riemann-Liouville fractional integrals*, Italian Journal of Pure and Applied Mathematics, **33**(2014), 299-306.
- [6] Chen, F., Wu, S., *Some Hermite-Hadamard type inequalities for harmonically s -convex functions*, The Scientific World Journal, Vol. 2014, Art. ID:279158, 2014.
- [7] Dragomir, S.S., *Refinements of the Hermite-Hadamard integral inequality for log-convex functions*, Aust. Math. Soc. Gaz., **28**(2001), no. 3, 129-134.
- [8] Hadamard, J., *Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann*, J. Math. Pures Appl., **58**(1983), 171-215.
- [9] İşcan, İ., *New general integral inequalities for quasi-geometrically convex functions via fractional integrals*, J. Inequal. Appl., **491**(2013), 15 pages.
- [10] İşcan, İ., *Hermite-Hadamard type inequalities for harmonically convex functions*, Hacet. J. Math. Stat., **43**(2014), no. 6, 935-942.
- [11] Kırmacı, U.S., Bakula, M.K., Özdemir, M.E., Pečarić, J., *Hadamard-type inequalities for s -convex functions*, Applied Mathematics and Computation, **193**(2007), no. 1, 26-35.
- [12] Kilbas, A.A., Srivastava, H.M., Trujillo, J.J., *Theory and applications of fractional differential equations*, Elsevier, Amsterdam, 2006.
- [13] Kunt, M., İşcan, İ., *On new inequalities of Hermite-Hadamard-Fejer type for GA-convex functions via fractional integrals*, RGMIA Research Report Collection, **18**(2015), Art. 108, 12 pp.
- [14] Niculescu, C.P., *Convexity according to the geometric mean*, Math. Inequal. Appl., **3**(2000), no. 2, 155-167.
- [15] Niculescu, C.P., *Convexity according to means*, Math. Inequal. Appl., **6**(2003), no. 4, 571-579.
- [16] Pachpatte, B.G., *On some inequalities for convex functions*, RGMIA Research Report Collection E, vol. 6, 2003.
- [17] Pachpatte, B.G., *A note on integral inequalities involving two log-convex functions*, Mathematical Inequalities and Applications, **7**(2004), no. 4, 511-515.
- [18] Rubinov, A.M., Dutta, J., *Hadamard inequality for quasi-convex functions in higher dimensions*, J. Math. Anal. Appl., **270**(2002), 80-91.
- [19] Sarikaya, M.Z., Sağlam, A., Yildırım, H., *On some Hadamard-type inequalities for h -convex functions*, Journal of Mathematical Inequalities, **2**(2008), no. 3, 335-341.
- [20] Yang, G.S., *Refinements of Hadamard inequality for r -convex functions*, Indian J. Pure Appl. Math., **32**(2001), no. 10, 1571-1579.
- [21] Yin, H.P., Qi, F., *Hermite-Hadamard type inequalities for the product of (α, m) -convex functions*, Journal of Nonlinear Science and Applications, **8**(2015), 231-236.

İmdat İşcan

Department of Mathematics

Faculty of Sciences and Arts

Giresun University

Giresun, Turkey

e-mail: imdat.iscan@giresun.edu.tr; imdati@yahoo.com

Mehmet Kunt

Department of Mathematics

Faculty of Sciences

Karadeniz Technical University

Trabzon, Turkey

e-mail: mkunt@ktu.edu.tr

