

On close-to-convex functions satisfying a differential inequality

Sukhwinder Singh Billing

Abstract. Let $\mathcal{C}_\alpha(\beta)$ denote the class of normalized functions f , analytic in the open unit disk \mathbb{E} which satisfy the condition

$$\Re \left[(1 - \alpha) \frac{zf'(z)}{\phi(z)} + \alpha \frac{(zf'(z))'}{\phi'(z)} \right] > \beta, \quad z \in \mathbb{E},$$

where $\frac{f(z)f'(z)}{z} \neq 0$, $z \in \mathbb{E}$, ϕ is starlike and α, β are pre-assigned real numbers. In 1977, Chichra, P. N. [1] introduced and studied the class $\mathcal{C}_\alpha = \mathcal{C}_\alpha(0)$. He proved the members of class \mathcal{C}_α are close-to-convex for $\alpha \geq 0$. We here prove that functions in class $\mathcal{C}_\alpha(\beta)$ are close-to-convex for $-\frac{\alpha}{2} \Re \left(\frac{z\phi'(z)}{\phi(z)} \right) \leq \beta < 1$, $\alpha \geq 0$ and the result is sharp in the sense that the constant β cannot be replaced by a real number smaller than $-\frac{\alpha}{2} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right)$. We claim that our result improves the result of Chichra, P. N. [1].

Mathematics Subject Classification (2010): 30C80, 30C45.

Keywords: Analytic function, convex function, starlike function, close-to-convex.

1. Introduction

Let \mathcal{A} be the class of functions f , analytic in $\mathbb{E} = \{z : |z| < 1\}$ and normalized by the conditions $f(0) = f'(0) - 1 = 0$. Let \mathcal{S}^* and \mathcal{K} denote the classes of starlike and convex functions respectively analytically defined as follows:

$$\mathcal{S}^* = \left\{ f \in \mathcal{A} : \Re \left(\frac{zf'(z)}{f(z)} \right) > 0, \quad z \in \mathbb{E} \right\},$$

and

$$\mathcal{K} = \left\{ f \in \mathcal{A} : \Re \left(1 + \frac{zf''(z)}{f'(z)} \right) > 0, \quad z \in \mathbb{E} \right\}.$$

This is well-known that

$$f(z) \in \mathcal{K} \Leftrightarrow zf'(z) \in \mathcal{S}^*. \tag{1.1}$$

A function $f \in \mathcal{A}$ is said to be close to convex if there is a real number $\alpha, -\pi/2 < \alpha < \pi/2$ and a convex function g (not necessarily normalized) such that

$$\Re \left(e^{i\alpha} \frac{f'(z)}{g'(z)} \right) > 0, \quad z \in \mathbb{E}.$$

In view of the relation (1.1), the above definition takes the following form in case g is normalized. A function $f \in \mathcal{A}$ is said to be close to convex if there is a real number $\alpha, -\pi/2 < \alpha < \pi/2$, and a starlike function ϕ such that

$$\Re \left(e^{i\alpha} \frac{zf'(z)}{\phi(z)} \right) > 0, \quad z \in \mathbb{E}.$$

It is well-known that every close-to-convex function is univalent. In 1934/35, Noshiro [3] and Warchawski [4] obtained a simple but elegant criterion for univalence of analytic functions. They proved that if an analytic function f satisfies $\Re f'(z) > 0$ for all z in \mathbb{E} , then f is close-to-convex and hence univalent in \mathbb{E} .

Let $\mathcal{C}_\alpha(\beta)$ denote the class of normalized analytic functions f which satisfy the condition

$$\Re \left[(1 - \alpha) \frac{zf'(z)}{\phi(z)} + \alpha \frac{(zf'(z))'}{\phi'(z)} \right] > \beta, \quad z \in \mathbb{E},$$

where $\frac{f(z)f'(z)}{\phi(z)} \neq 0, z \in \mathbb{E}$, ϕ is starlike and α, β are pre-assigned real numbers. The class $\tilde{\mathcal{C}}_\alpha = \mathcal{C}_\alpha(0)$ was introduced and studied by Chichra, P. N. [1] in 1977. He called the members of class \mathcal{C}_α as α -close-to-convex functions. Infact, he proved the following result.

Theorem 1.1. *Let $f(z) \in \mathcal{C}_\alpha$ and $\alpha \geq 0$. Then $f(z)$ is close-to-convex in \mathbb{E} .*

In the present paper, we establish the result that functions in $\mathcal{C}_\alpha(\beta)$ are close-to-convex for $-\frac{\alpha}{2} \Re \left(\frac{z\phi'(z)}{\phi(z)} \right) \leq \beta < 1, \alpha \geq 0$. Our result is the best possible in the sense that

the constant β cannot be replaced by a real number smaller than $-\frac{\alpha}{2} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right)$.

We also claim that our result improves the result of Chichra, P. N. [1]. To prove our main result, we shall use the following lemma of Miller [2].

Lemma 1.2. *Let \mathbb{D} be a subset of $\mathbb{C} \times \mathbb{C}$ (\mathbb{C} is the complex plane) and let $\phi : \mathbb{D} \rightarrow \mathbb{C}$ be a complex function. For $u = u_1 + iu_2, v = v_1 + iv_2$ (u_1, u_2, v_1, v_2 are reals), let ϕ satisfy the following conditions:*

- (i) $\phi(u, v)$ is continuous in \mathbb{D}
- (ii) $(1, 0) \in \mathbb{D}$ and $\Re[\phi(1, 0)] > 0$ and
- (iii) $\Re[\phi(iu_2, v_1)] \leq 0$ for all $(iu_2, v_1) \in \mathbb{D}$ such that $v_1 \leq -(1 + u_2^2)/2$.

Let $p(z) = 1 + p_1z + p_2z^2 + \dots$ be regular in the open unit disk \mathbb{E} , such that $(p(z), zp'(z)) \in \mathbb{D}$ for all $z \in \mathbb{E}$. If

$$\Re[\phi(p(z), zp'(z))] > 0, \quad z \in \mathbb{E},$$

then $\Re p(z) > 0, z \in \mathbb{E}$.

2. Main result

Theorem 2.1. *Let α and β be real numbers such that $\alpha \geq 0$ and*

$$-\frac{\alpha}{2} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right) \leq \beta < 1$$

for a starlike function ϕ . Assume that $f \in \mathcal{A}$ satisfies

$$\Re \left[(1 - \alpha) \frac{zf'(z)}{\phi(z)} + \alpha \frac{(zf'(z))'}{\phi'(z)} \right] > \beta, \quad z \in \mathbb{E}, \tag{2.1}$$

then $\Re \left(\frac{zf'(z)}{\phi(z)} \right) > 0$ in \mathbb{E} and hence f is close-to-convex and hence univalent in \mathbb{E} . The result is sharp in the sense that the constant β on the right hand side of (2.1) cannot be replaced by a real number smaller than $-\frac{\alpha}{2} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right)$.

Proof. Let $p(z) = 1 + p_1z + p_2z^2 + \dots$ be analytic in \mathbb{E} such that for all $z \in \mathbb{E}$, we write

$$\frac{zf'(z)}{\phi(z)} = p(z).$$

Then,

$$(1 - \alpha) \frac{zf'(z)}{\phi(z)} + \alpha \frac{(zf'(z))'}{\phi'(z)} = p(z) + \alpha zp'(z) \frac{\phi(z)}{z\phi(z)}.$$

Therefore, condition (2.1) is equivalent to

$$\Re \left(\frac{1}{1 - \beta} p(z) + \frac{\alpha}{1 - \beta} zp'(z) \frac{\phi(z)}{z\phi'(z)} - \frac{\beta}{1 - \beta} \right) > 0, \quad z \in \mathbb{E}. \tag{2.2}$$

For $\mathbb{D} = \mathbb{C} \times \mathbb{C}$, define $\Phi(u, v) : \mathbb{D} \rightarrow \mathbb{C}$ as under:

$$\Phi(u, v) = \frac{1}{1 - \beta} u + \frac{\alpha}{1 - \beta} v \frac{\phi(z)}{z\phi'(z)} - \frac{\beta}{1 - \beta}, \quad z \in \mathbb{E}.$$

Then $\Phi(u, v)$ is continuous in \mathbb{D} , $(1, 0) \in \mathbb{D}$ and $\Re(\Phi(1, 0)) = 1 > 0$. Further, in view of (2.2), we get, $\Re[\Phi(p(z), zp'(z))] > 0, z \in \mathbb{E}$. Let $u = u_1 + iu_2, v = v_1 + iv_2$ where u_1, u_2, v_1 and v_2 are all real numbers. Then, for $(iu_2, v_1) \in \mathbb{D}$, with $v_1 \leq -\frac{1 + u_2^2}{2}$, we have

$$\begin{aligned} \Re \Phi(iu_2, v_1) &= \Re \left(\frac{1}{1 - \beta} u_2 i + \frac{\alpha}{1 - \beta} v_1 \frac{\phi(z)}{z\phi'(z)} - \frac{\beta}{1 - \beta} \right) \\ &\leq - \left[\frac{\alpha}{1 - \beta} \frac{1 + u_2^2}{2} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right) + \frac{\beta}{1 - \beta} \right] \\ &\leq - \left[\frac{\alpha}{2(1 - \beta)} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right) + \frac{\beta}{1 - \beta} \right] \\ &\leq 0. \end{aligned}$$

In view of (2.2) and Lemma 1.2, proof now follows.




Figure 2.1




Figure 2.2

To show that the constant β on the right hand side of (2.1) cannot be replaced by a real number smaller than $-\frac{\alpha}{2} \Re \left(\frac{\phi(z)}{z\phi'(z)} \right)$, we consider the function $f(z) = z e^z \in \mathcal{A}$ and $\phi(z) = z \in \mathcal{S}^*$. Using Mathematica 9.0, we plot, in Figure 2.1, the image of the unit disk under the operator $(1 - \alpha) \frac{zf'(z)}{\phi(z)} + \alpha \frac{(zf'(z))'}{\phi'(z)}$ taking $\alpha = 2$. From this figure, we notice that minimum real part of $(1 - \alpha) \frac{zf'(z)}{\phi(z)} + \alpha \frac{(zf'(z))'}{\phi'(z)}$ is smaller

than -1 (the calculated value of $-\frac{\alpha}{2}\Re\left(\frac{\phi(z)}{z\phi'(z)}\right)$ for $\alpha = 2$ and $\phi(z) = z$). In Figure 2.2, we plot the image of unit disk under the operator $\frac{zf'(z)}{\phi(z)}$. It is obvious that $\Re\left(\frac{zf'(z)}{\phi(z)}\right) \not\geq 0$ for all z in \mathbb{E} . For example, the point $z = -\frac{1}{2} + i\frac{\pi}{4}$ is an interior point of \mathbb{E} , but at this point $\Re\left(\frac{zf'(z)}{\phi(z)}\right) = -\frac{\pi - 2}{4\sqrt{2}e} = -0.1224\dots < 0$. This justifies our claim. □

Remark 2.2. We claim that our result improves the result of Chichra, P. N. [1]. In fact, when we take $f(z) = -z - 2\log(1 - z) \in \mathcal{A}$, $\phi(z) = z$ and $\alpha = 2$ in Theorem 2.1, we notice that at $z = -1$,

$$\Re\left[(1 - \alpha)\frac{zf'(z)}{\phi(z)} + \alpha\frac{(zf'(z))'}{\phi'(z)}\right] = -1.$$

Thus the function f does not satisfy the hypothesis of Theorem 1.1 due to Chichra, P. N. [1] i.e. $f \notin \mathcal{C}_\alpha$ although $\Re\left(\frac{zf'(z)}{\phi(z)}\right) = \Re\left(\frac{1+z}{1-z}\right) > 0$ in \mathbb{E} . Hence the result of Chichra, P. N. [1] fails to conclude the close-to-convexity in this case whereas Theorem 2.1 concludes the same.

References

- [1] Chichra, P.N., *New subclasses of the class of close-to-convex functions*, Proc. Amer. Math. Soc., **62**(1)(1977), 37-43.
- [2] Miller, S.S., *Differential inequalities and Carathéodory functions*, Bull. Amer. Math. Soc., **81**(1975), 79-81.
- [3] Noshiro, K., *On the theory of schlicht functions*, J. Fac. Sci., Hokkaido Univ., **2**(1934-35), 129-155.
- [4] Warchawski, S.E., *On the higher derivatives at the boundary in conformal mappings*, Trans. Amer. Math. Soc., **38**(1935), 310-340.

Sukhwinder Singh Billing
 Department of Mathematics
 Sri Guru Granth Sahib World University
 Fatehgarh Sahib-140 406, Punjab, India
 e-mail: ssbilling@gmail.com

