

# On applications of Andrica-Badea and Nagy inequalities in spectral graph theory

Igor Milovanović, Emina Milovanović and Edin Glogić

**Abstract.** Applications of Andrica-Badea and Nagy inequalities for determining bounds of graph invariants of undirected, connected graphs are investigated. We consider bounds of the following invariants: the first Zagreb index, general Randić index, Laplacian linear spread and normalized Laplacian spread of graphs.

**Mathematics Subject Classification (2010):** 60E15, 05C50.

**Keywords:** Inequalities, first Zagreb index, spread of graph.

## 1. Introduction

Andrica and Badea (see [1]) have proved the following result.

Let  $p_1, p_2, \dots, p_n$  be non-negative real numbers and  $a_1, a_2, \dots, a_n$  and  $b_1, b_2, \dots, b_n$  real numbers with the properties

$$0 < r_1 \leq a_i \leq R_1 < +\infty \quad \text{and} \quad 0 < r_2 \leq b_i \leq R_2 < +\infty$$

for each  $i = 1, 2, \dots, n$ . Further, let  $S$  be a subset of  $I_n = \{1, 2, \dots, n\}$  which minimizes the expression

$$\left| \sum_{i \in S} p_i - \frac{1}{2} \sum_{i=1}^n p_i \right|. \quad (1.1)$$

Then

$$\begin{aligned} & \left| \sum_{i=1}^n p_i \sum_{i=1}^n p_i a_i b_i - \sum_{i=1}^n p_i a_i \sum_{i=1}^n p_i b_i \right| \\ & \leq (R_1 - r_1)(R_2 - r_2) \sum_{i \in S} p_i \left( \sum_{i=1}^n p_i - \sum_{i \in S} p_i \right) \end{aligned} \quad (1.2)$$

In [17] Nagy has proved the following result:

Let  $a_1, a_2, \dots, a_n$  are real numbers with the property  $r \leq a_i \leq R$ , for each  $i = 1, 2, \dots, n$ . Then

$$n \sum_{i=1}^n a_i^2 - \left( \sum_{i=1}^n a_i \right)^2 \geq \frac{n}{2} (R - r)^2. \quad (1.3)$$

In this paper we consider bounds of some graph invariants and prove that they are direct corollaries of inequalities (1.2) and (1.3). Some of the obtained bounds are better than those obtained in the literature so far.

In the next section we recall some results from spectral graph theory needed for our work.

## 2. Laplacian and normalized laplacian spectrum of graph

Let  $G = (V, E)$ ,  $V = \{1, 2, \dots, n\}$ , be undirected connected graph with  $n$  vertices and  $m$  edges, with sequence of vertex degrees  $d_1 \geq d_2 \geq \dots \geq d_n > 0$ ,  $d_i = d(i)$ ,  $i = 1, 2, \dots, n$ . Denote with  $\mathbf{A}$  adjacency matrix of  $G$ . Its eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$  represent ordinary eigenvalues of graph  $G$ . If  $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n)$  is diagonal matrix of vertex degrees, then  $\mathbf{L} = \mathbf{D} - \mathbf{A}$  is the Laplacian matrix of the  $G$ . Eigenvalues of  $\mathbf{L}$ ,  $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} > \mu_n = 0$  represent Laplacian eigenvalues of graph  $G$ . The main properties of these eigenvalues are (see [3, 7, 8, 15])

$$\sum_{i=1}^{n-1} \mu_i = \sum_{i=1}^n d_i = 2m \quad \text{and} \quad \sum_{i=1}^{n-1} \mu_i^2 = \sum_{i=1}^n d_i^2 + \sum_{i=1}^n d_i = M_1 + 2m, \quad (2.1)$$

where  $M_1 = \sum_{i=1}^n d_i^2$  is the first Zagreb index (see [13]).

Because the graph  $G$  is assumed to be connected, it has no isolated vertices and therefore the matrix  $\mathbf{D}^{-1/2}$  is well-defined. Then  $\mathbf{L}^* = \mathbf{D}^{-1/2} \mathbf{L} \mathbf{D}^{-1/2}$  is the normalized Laplacian matrix of the graph  $G$ . Its eigenvalues are  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_{n-1} > \rho_n = 0$ . Main properties of these eigenvalues are given by (see [19])

$$\sum_{i=1}^{n-1} \rho_i = n \quad \text{and} \quad \sum_{i=1}^{n-1} \rho_i^2 = n + 2R_{-1}, \quad (2.2)$$

where  $R_{-1} = \sum_{\{i,j\} \in E} (d_i d_j)^{-1}$  is the general Randić index (see [6, 18]).

## 3. Main result

In the following theorem we prove the inequality that establishes lower and upper bounds for invariant  $M_1$  in terms of parameters  $n$ ,  $m$ ,  $d_1$  and  $d_n$ .

**Theorem 3.1.** *Let  $G = (V, E)$  be undirected connected graph with  $n, n \geq 2$ , vertices and  $m$  edges. Then*

$$\frac{4m^2}{n} + \frac{1}{2}(d_1 - d_n)^2 \leq M_1 \leq \frac{4m^2}{n} + n\alpha(n)(d_1 - d_n)^2, \quad (3.1)$$

where

$$\alpha(n) = \frac{1}{n} \left\lfloor \frac{n}{2} \right\rfloor \left( 1 - \frac{1}{n} \left\lfloor \frac{n}{2} \right\rfloor \right) = \frac{1}{4} \left( 1 - \frac{(-1)^{n+1} + 1}{2n^2} \right) = \begin{cases} \frac{1}{4}, & \text{if } n \text{ is even} \\ \frac{n^2-1}{4n^2}, & \text{if } n \text{ is odd} \end{cases}$$

Equality holds if and only if  $G$  is regular graph.

*Proof.* For  $a_i = d_i$ ,  $i = 1, 2, \dots, n$ ,  $R = d_1$  and  $r = d_n$ , the inequality (1.3) transforms into

$$n \sum_{i=1}^n d_i^2 - \left( \sum_{i=1}^n d_i \right)^2 \geq \frac{n}{2} (d_1 - d_n)^2,$$

i.e. according to (2.1), into

$$nM_1 - 4m^2 \geq \frac{n}{2} (d_1 - d_n)^2, \quad (3.2)$$

wherefrom the left part of inequality (3.1) is obtained.

For  $p_i = 1$ ,  $i = 1, 2, \dots, n$  and  $S = \{1, 2, \dots, k\} \subset I_n$ , the expression (1.1) reaches the minimum for  $k = \left\lfloor \frac{n}{2} \right\rfloor$ . Now for  $S = \{1, 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor\}$ ,  $p_i = 1$ ,  $a_i = b_i = d_i$ ,  $i = 1, 2, \dots, n$ ,  $r_1 = r_2 = d_n$  and  $R_1 = R_2 = d_1$ , the inequality (1.2) becomes

$$n \sum_{i=1}^n d_i^2 - \left( \sum_{i=1}^n d_i \right)^2 \leq (d_1 - d_n)^2 \left\lfloor \frac{n}{2} \right\rfloor \left( n - \left\lfloor \frac{n}{2} \right\rfloor \right)$$

i.e.

$$nM_1 - 4m^2 \leq n^2 (d_1 - d_n)^2 \alpha(n) \quad (3.3)$$

where

$$\alpha(n) = \frac{1}{n} \left\lfloor \frac{n}{2} \right\rfloor \left( 1 - \frac{1}{n} \left\lfloor \frac{n}{2} \right\rfloor \right) = \frac{1}{4} \left( 1 - \frac{(-1)^{n+1} + 1}{2n^2} \right) = \begin{cases} \frac{1}{4}, & \text{if } n \text{ is even} \\ \frac{n^2-1}{4n^2}, & \text{if } n \text{ is odd} \end{cases}.$$

From (3.3) right side of inequality (3.1) immediately follows.

Equalities in (3.2) and (3.3) hold if and only if  $d_1 = d_2 = \dots = d_n$ , so the equalities in (3.1) hold if and only if  $G$  is a regular graph.  $\square$

**Remark 3.2.** Since  $(d_1 - d_n)^2 \geq 0$ , left inequality in (3.1) is stronger than

$$M_1 \geq \frac{4m^2}{n} \quad (3.4)$$

which was proved in [9].

**Remark 3.3.** In [2] the invariant  $irr_{EB}(G) = \sqrt{\frac{nM_1}{4m^2} - 1}$  as the irregularity measure of graph was introduced. In [11] another irregularity measure  $irr_g(G) = \frac{d_1}{d_n} - 1$  was defined. According to (3.1) we can establish the following relationship between these two measures

$$\sqrt{\frac{nd_n}{8m^2}} irr_g(G) \leq irr_{EB}(G) \leq \sqrt{\frac{n^2 d_n \alpha(n)}{4m^2}} irr_g(G).$$

The linear Laplacian spread of graph  $G$  is defined as  $\mu_1 - \mu_{n-1}$ . The following theorem establishes lower and upper bounds for this invariant in terms of parameters  $n$ ,  $m$  and  $M_1$ .

**Theorem 3.4.** *Let  $G = (V, E)$  be undirected connected graph with  $n, n \geq 2$ , vertices and  $m$  edges. Then*

$$\sqrt{\frac{(n-1)(M_1+2m)-4m^2}{(n-1)^2\alpha(n-1)}} \leq \mu_1 - \mu_{n-1} \leq \sqrt{\frac{2((n-1)(M_1+2m)-4m^2)}{n-1}}. \quad (3.5)$$

where

$$\alpha(n-1) = \frac{1}{n-1} \left\lfloor \frac{n-1}{2} \right\rfloor \left( 1 - \frac{1}{n-1} \left\lfloor \frac{n-1}{2} \right\rfloor \right) = \frac{1}{4} \left( 1 - \frac{(-1)^n + 1}{2(n-1)^2} \right).$$

Equalities hold if and only if  $G$  is a complete graph,  $G \cong K_n$ .

*Proof.* For  $n := n-1$ ,  $a_i = \mu_i$ ,  $i = 1, 2, \dots, n-1$ ,  $R = \mu_1$  and  $r = \mu_{n-1}$ , the inequality (1.3) becomes

$$(n-1) \sum_{i=1}^{n-1} \mu_i^2 - \left( \sum_{i=1}^{n-1} \mu_i \right)^2 \geq \frac{(n-1)}{2} (\mu_1 - \mu_{n-1})^2$$

i.e. according to (2.1) we have

$$(n-1)(M_1+2m)-4m^2 \geq \frac{(n-1)}{2} (\mu_1 - \mu_{n-1})^2 \quad (3.6)$$

wherefrom right side of (3.5) is obtained.

For  $n := n-1$ ,  $p_i = 1$ ,  $a_i = b_i = \mu_i$ ,  $i = 1, 2, \dots, n-1$ ,  $r_1 = r_2 = \mu_{n-1}$  and  $R_1 = R_2 = \mu_1$ , the inequality (1.2) transforms into

$$(n-1) \sum_{i=1}^{n-1} \mu_i^2 - \left( \sum_{i=1}^{n-1} \mu_i \right)^2 \leq (\mu_1 - \mu_{n-1})^2 \left\lfloor \frac{n-1}{2} \right\rfloor \left( n-1 - \left\lfloor \frac{n-1}{2} \right\rfloor \right)$$

i.e.

$$(n-1)(M_1+2m)-4m^2 \leq (\mu_1 - \mu_{n-1})^2 (n-1)^2 \alpha(n), \quad (3.7)$$

where

$$\alpha(n-1) = \frac{1}{n-1} \left\lfloor \frac{n-1}{2} \right\rfloor \left( 1 - \frac{1}{n-1} \left\lfloor \frac{n-1}{2} \right\rfloor \right) = \begin{cases} \frac{1}{4}, & \text{if } n \text{ is even} \\ \frac{n(n-2)}{4(n-1)^2}, & \text{if } n \text{ is odd} \end{cases}.$$

From (3.7) left part of inequality (3.5) is directly obtained.

Equalities (3.6) and (3.7) hold if and only if  $\mu_1 = \mu_2 = \dots = \mu_{n-1}$ , hence equalities in (3.5) hold if and only if  $G$  is a complete graph,  $G \cong K_n$ .  $\square$

**Remark 3.5.** Right side of inequality (3.5) was proved in [16]. Since  $\alpha(n-1) \leq \frac{1}{4}$ , for each  $n$ , left side of inequality (3.5) is stronger than

$$\mu_1 - \mu_{n-1} \geq \frac{2}{n-1} \sqrt{(n-1)(M_1+2m)-4m^2},$$

for even  $n$ . The above inequality was proved in [10] and [20].

From inequality (3.4) and  $M_1 \leq m \left( \frac{2m}{n-1} + (n-2) \right)$ , proved in [5], and inequality  $M_1 \geq \frac{4m^2}{n}$ , proved in [9], the following corollary of Theorem 3.4 holds.

**Corollary 3.6.** *Let  $G = (V, E)$  be undirected connected graph with  $n, n \geq 2$ , vertices and  $m$  edges. Then*

$$\frac{1}{n-1} \sqrt{\frac{2m(n(n-1)-2m)}{n\alpha(n-1)}} \leq \mu_1 - \mu_{n-1} \leq \sqrt{\frac{2m(n(n-1)-2m)}{n-1}}.$$

*Equalities hold if and only if  $G$  is a complete graph,  $G \cong K_n$ .*

In the following theorem we determine lower and upper bounds for graph invariant  $R_{-1}$  in terms of parameters  $n$ ,  $\rho_1$  and  $\rho_{n-1}$ .

**Theorem 3.7.** *Let  $G = (V, E)$  be undirected connected graph with  $n, n \geq 2$ , vertices and  $m$  edges. Then*

$$\frac{n}{2(n-1)} + \frac{1}{4}(\rho_1 - \rho_{n-1})^2 \leq R_{-1} \leq \frac{n}{2(n-1)} + \frac{(n-1)\alpha(n-1)}{2}(\rho_1 - \rho_{n-1})^2. \quad (3.8)$$

*Equalities hold if and only if  $G$  is a complete graph,  $G \cong K_n$ .*

*Proof.* for  $n := n-1$ ,  $a_i = \rho_i$ ,  $i = 1, 2, \dots, n-1$ ,  $r = \rho_{n-1}$  and  $R = \rho_1$  the inequality (1.3) becomes

$$(n-1) \sum_{i=1}^{n-1} \rho_i^2 - \left( \sum_{i=1}^{n-1} \rho_i \right)^2 \geq \frac{n-1}{2}(\rho_1 - \rho_{n-1})^2$$

i.e. according to (2.2)

$$(n-1)(n+2R_{-1}) - n^2 \geq \frac{n-1}{2}(\rho_1 - \rho_{n-1})^2, \quad (3.9)$$

wherefrom left side of inequality (3.8) is obtained.

For  $n := n-1$ ,  $p_i = 1$ ,  $a_i = b_i = \rho_i$ ,  $i = 1, 2, \dots, n-1$ ,  $r_1 = r_2 = \rho_{n-1}$  and  $R_1 = R_2 = \rho_1$ , inequality (1.3) transforms into

$$(n-1) \sum_{i=1}^{n-1} \rho_i^2 - \left( \sum_{i=1}^{n-1} \rho_i \right)^2 \leq (\rho_1 - \rho_{n-1})^2 \left\lfloor \frac{n-1}{2} \right\rfloor \left( n-1 - \left\lfloor \frac{n-1}{2} \right\rfloor \right),$$

i.e.

$$(n-1)(n+2R_{-1}) - n^2 \leq (n-1)^2 \alpha(n-1) (\rho_1 - \rho_{n-1})^2, \quad (3.10)$$

wherefrom right part of inequality (3.8) is obtained.

Equalities in (3.9) and (3.10) hold if and only if  $\rho_1 = \rho_2 = \dots = \rho_{n-1}$ , therefore equalities in (3.8) hold if and only if  $G$  is complete graph,  $G \cong K_n$ .  $\square$

**Remark 3.8.** Since  $(\rho_1 - \rho_{n-1})^2 \geq 0$ , it follows that left side of inequality (3.8) is stronger than inequality

$$R_{-1} \geq \frac{n}{2(n-1)}$$

which was proved in [14].

**Remark 3.9.** Inequalities in (3.8) can be presented in an equivalent form as

$$\sqrt{\frac{2(n-1)R_{-1}-n}{(n-1)^2\alpha(n-1)}} \leq \rho_1 - \rho_{n-1} \leq \sqrt{\frac{2(2(n-1)R_{-1}-n)}{n-1}}. \quad (3.11)$$

For even  $n$ , left side of inequality (3.11) is stronger than inequality

$$\rho_1 - \rho_{n-1} \geq \frac{2}{n-1} \sqrt{2(n-1)R_{-1}-n},$$

which was proved in [4].

**Acknowledgement.** This work was supported by the Serbian Ministry of Education and Science, Project No TR32012 and TR32009.

## References

- [1] Andrica, D., Badea, C., *Grös inequality for positive linear functionals*, Period. Math. Hungar, **19**(1988), no. 2, 155–167.
- [2] Bell, F.K., *A note on the irregularity of graphs*, Lin. Algebra Appl., **161**(1992), 45–54.
- [3] Biggs, N., *Algebraic graph theory*, Cambridge Univ. Press, Cambridge, UK, 1993.
- [4] Butler, S., *Eigenvalues and structures of graph*, Ph. Dissertation, University of California, San Diego, 2008.
- [5] de Caen, D., *An upper bound on the sum of squares of degrees in a graph*, Discr. Math., **185**(1998), no. 1-3, 245–248.
- [6] Cavers, M., Fallat, S., Kirkland, S., *On the normalized Laplacian energy and general Randic index of graphs*, Lin. Algebra Appl., (2010), no. 433, 172–190.
- [7] Chung, F.R.K., *Spectral graph theory*, Am. Math. Soc., Providence, 1997.
- [8] Cvetković, D., Doob, M., Sachs, H., *Spectra of graph theory and application*, New York, Academic Press, 1980.
- [9] Edwards, C.S., *The largest vertex degree sum for a triangle of a graph*, Bull. London Math. Soc., **9**(1977), 203–208.
- [10] Fath-Tabar, G.H., Ashrafi, A.R., *Some remark on Laplacian eigenvalues of graphs*, Math. Commun., **15**(2010), no. 2, 443–451.
- [11] Goldberg, F., *New results on eigenvalues on degree deviation* <http://orxiv.org/1403.269V1>, 2014.
- [12] Gutman, I., Furtula, B., Elphick, C., *Tree new/old vertex-degree based topological indices*, MATCH Commun. Math. Comput. Chem., **72**(2014), 617–632.
- [13] Gutman, I., Trinajstić, N., *Graph theory and molecular orbitals. Total  $\pi$ -electron energy of alternant hydrocarbons*, Chem. Phys. Letters, **17**(1972), 535–538.
- [14] Li, X., Yang, Y., *Sharp bounds for the general Randić index*, MATCH Commun. Math. Comput. Chem., **51**(2004), 155–166.
- [15] Merris, R., *Laplacian matrices of graphs: A survey*, Lin. Algebra Appl., **197-198**(1994), 143–176.
- [16] Milovanović, I., Milovanović, E., *Remark on inequalities for the Laplacian spread of graphs*, Czech. Math. J., **64** (2014), no. 139, 285–287.

- [17] Nagy, J.V.S., *Über algebraische Gleichungen mit lauter reellen Wurzeln*, Jahresbericht der deutschen mathematiker-Vereinigung, **27**(1918), 37-43.
- [18] Randić, M., *On characterization of molecular branching*, J. Amer. Chem. Soc., **97**(1975), 6609–6615.
- [19] Zumstein, P., *Comparison of spectral methods through the adjacency matrix and the Laplacian of a graph*, Ph. Dissertation, ETH Zürich, 2005.
- [20] You, Z., Liu, B., *The Laplacian spread of graphs*, Czech. Math. J., **62**(2012), 159–168.

Igor Milovanović  
 Faculty of Electronic Engineering  
 Beogradska 14, 18000 Niš, Serbia  
 e-mail: [igor@elfak.ni.ac.rs](mailto:igor@elfak.ni.ac.rs)

Emina Milovanović  
 Faculty of Electronic Engineering  
 Beogradska 14, 18000 Niš, Serbia  
 e-mail: [ema@elfak.ni.ac.rs](mailto:ema@elfak.ni.ac.rs)

Edin Glogić  
 State University of Novi Pazar  
 Novi Pazar, Serbia  
 e-mail: [edin\\_gl@hotmail.com](mailto:edin_gl@hotmail.com)

