

New integral inequalities involving generalized Riemann-Liouville fractional operators

Juan Gabriel Galeano Delgado,
Juan E. Nápoles Valdés and Edgardo Pérez Reyes

Abstract. In this paper, using a generalized operator of the Riemann-Liouville type, defined and studied in a previous work, several integral inequalities for synchronous functions are established.

Mathematics Subject Classification (2010): 26A33, 26D10, 47A63.

Keywords: Generalized fractional Riemann-Liouville integral, fractional integral inequality, synchronous functions.

1. Introduction


One of the most developed mathematical areas in the last 20 years is that of Integral Inequalities, associated with different functional notions: convex, synchronous functions among other, within the framework of Riemann, fractional and generalized integral operators. A detail that we want to point out is the fact of the appearance in recent years of various integral operators, natural extensions of the fractional integral of Riemann-Liouville, this together with the attention received by Integral Inequalities, make more and more researchers and research is devoted to this topic. To get a more complete idea in this regard, we recommend consulting the works [1, 2, 6, 8, 10, 11, 12, 13, 16] and the references cited therein.

In this direction, one of the most fruitful notions is the following (see [4]).

Definition 1.1. If χ and ψ are two integrable functions on $[a, b]$, they are synchronous on $[a_1, a_2]$ if $(\chi(x) - \chi(y))(\psi(x) - \psi(y)) \geq 0$, for any $x, y \in [a_1, a_2]$.

Received 12 July 2020; Accepted 13 August 2020.

© Studia UBB MATHEMATICA. Published by Babeş-Bolyai University

 This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

Various integral inequalities have been obtained using this notion, within the framework of different integral operators (see [4], [7], [9], [15], [18], [20], and the references cited there).

Throughout the work we use the functions Γ (see [17, 19, 22, 23]) and Γ_k (cf. defined by [5]):

$$\Gamma(z) = \int_0^\infty \tau^{z-1} e^{-\tau} d\tau, \quad \Re(z) > 0, \tag{1.1}$$

$$\Gamma_k(z) = \int_0^\infty \tau^{z-1} e^{-\tau^k/k} d\tau, k > 0. \tag{1.2}$$

It is clear that if $k \rightarrow 1$ we have $\Gamma_k(z) \rightarrow \Gamma(z)$, $\Gamma_k(z) = (k)^{\frac{z}{k}-1} \Gamma(\frac{z}{k})$ and $\Gamma_k(z+k) = z\Gamma_k(z)$. As well, we define the k -beta function as follows

$$B_k(u, v) = \frac{1}{k} \int_0^1 \tau^{\frac{u}{k}-1} (1-\tau)^{\frac{v}{k}-1} d\tau,$$

notice that $B_k(u, v) = \frac{1}{k} B(\frac{u}{k}, \frac{v}{k})$ and $B_k(u, v) = \frac{\Gamma_k(u)\Gamma_k(v)}{\Gamma_k(u+v)}$.

In [7] the following fractional integral operator of the Riemann-Liouville type is defined.

Definition 1.2. The k -generalized fractional Riemann-Liouville integral of order α with $\alpha \in \mathbb{R}$, and $s \neq -1$ of an integrable function $\chi(u)$ on $[0, \infty)$, are given as follows:

$${}_s J_{F, a_1}^{\frac{\alpha}{k}} \chi(u) = \frac{1}{k\Gamma_k(\alpha)} \int_{a_1}^u \frac{F(\tau, s)\chi(\tau)d\tau}{[\mathbb{F}(u, \tau)]^{1-\frac{\alpha}{k}}}, \tag{1.3}$$

with $F(\tau, 0) = 1$ and $\mathbb{F}(u, \tau) = \int_\tau^u F(\theta, s)d\theta$.

Remark 1.3. In the aforementioned paper, the main properties of this operator (boundedness, commutativity, etc.) and various inequalities associated with it were studied.

Remark 1.4. If in Definition 1.2 we consider the kernel $F(t, s) = 1$ and $k = 1$, we obtain the classic fractional Integral Riemann-Liouville, used in the work [4]; in the case of the same kernel but $k \neq 1$ then the k -fractional integral of the Riemann-Liouville type of [14] is obtained (see also [15, 18, 20]). If, on the contrary, we consider the kernel $F(t, s) = t^s$, we obtain the $(k;s)$ -Riemann-Liouville fractional integral of [21]. In the case of taking the kernel as $F(t, s) = h'(t)$, we obtain the $(k;h)$ -Riemann-Liouville integral fractional used in [9]. It is clear then, that the results obtained in our work, generalize those of the works mentioned before.

The main purpose of this paper, using the generalized fractional integral operator of the Riemann-Liouville type of Definition 1.2, is to establish several integral inequalities, which contain as particular cases, several of those reported in the literature.

2. Main results

Our first fundamental result is the following.

Theorem 2.1. *Let φ, ψ be two synchronous functions on $[0, \infty)$ and let $f, \phi, \omega \geq 0$, then for all $\tau > a_1 \geq 0$, $\alpha > 0$, and $s \neq -1$, we have*

$$\begin{aligned}
 & 2 {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi\psi)(\tau) \right] \\
 & \quad + 2 {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi\psi)(\tau) \\
 & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\psi)(\tau) \right] \\
 & \quad + {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right] \\
 & \quad + {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right]. \tag{2.1}
 \end{aligned}$$

To prove the previous Theorem, we need the following lemma.

Lemma 2.2. *Let φ, ψ be two synchronous functions on $[0, \infty)$ and let $h, g \geq 0$, then for all $\tau > a_1 \geq 0$, $\alpha > 0$, and $s \neq -1$, we have the following inequality*

$$\begin{aligned}
 & {}^s J_{F,a_1}^{\frac{\alpha}{k}} h(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (g\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} g(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi\psi)(\tau) \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (g\psi)(\tau) \\
 & \quad + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (g\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\psi)(\tau). \tag{2.2}
 \end{aligned}$$

Proof. Since φ, ψ are synchronous functions on $[0, \infty)$, then for all $u, v \geq 0$, we have

$$\varphi(u)\psi(u) + \varphi(v)\psi(v) \geq \varphi(u)\psi(v) + \varphi(v)\psi(u). \tag{2.3}$$

Thus, if we multiple both sides of (2.3) by $\frac{F(u,s)h(u)}{k\Gamma_k(\alpha)[\mathbb{F}(\tau,u)]^{1-\frac{\alpha}{k}}}$, and then we integrate the resulting inequality with respect to u over (a_1, τ) , it holds that

$${}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi\psi)(\tau) + \varphi(v)\psi(v) {}^s J_{F,a_1}^{\frac{\alpha}{k}} h(\tau) \geq \psi(v) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi)(\tau) + \varphi(v) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\psi)(\tau). \tag{2.4}$$

Now, multiplying both sides of (2.4) by $\frac{F(v,s)g(v)}{k\Gamma_k(\alpha)[\mathbb{F}(\tau,v)]^{1-\frac{\alpha}{k}}}$, then we integrate the resulting inequality with respect to v over (a_1, τ) , we get

$$\begin{aligned}
 & {}^s J_{F,a_1}^{\frac{\alpha}{k}} h(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (g\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} g(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi\psi)(\tau) \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (g\psi)(\tau) \\
 & \quad + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (g\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\psi)(\tau).
 \end{aligned}$$

Thus, we conclude the result. □

Remark 2.3. If we consider the kernel $F(t, s) = 1$ and $k = 1$ of this result, we obtain the Lemma 3 of [4], in the case that $F(t, s) = h'(t)$, this result covers Lemma 1 of [9].

Let us now prove the Theorem 2.1.

Proof. By setting $h = f$ and $g = \phi$ in (2.2), then we multiple the resulting inequality by ${}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau)$, we get

$$\begin{aligned} & {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi\psi)(\tau) \right] \\ & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\psi)(\tau) \right]. \end{aligned} \tag{2.5}$$

By setting $h = \omega$ and $g = \phi$ in (2.2), and multiplying the result by ${}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau)$, we get

$$\begin{aligned} & {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi\psi)(\tau) \right] \\ & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right]. \end{aligned} \tag{2.6}$$

Now, using the same idea, we put $h = \omega$ and $g = f$, and then we multiple the result by ${}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau)$, we find

$$\begin{aligned} & {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi\psi)(\tau) \right] \\ & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right]. \end{aligned} \tag{2.7}$$

Finally, by adding the inequalities (2.5), (2.6) and (2.7) we obtain the result. □

Remark 2.4. If we take the kernel $F(t, s) = 1$ and $k = 1$ we obtain Theorem 2 of [4], and if $F(t, s) = h'(t)$, this result reduces to Theorem 3 of [9].

The next Lemma will be from very useful for proving the last theorem.

Lemma 2.5. *Let φ, ψ be two synchronous functions on $[0, \infty)$ and let $h, g \geq 0$, then for all $\tau > a_1 \geq 0$, $\alpha, \beta > 0$, and $s \neq -1$, we have the following inequality*

$$\begin{aligned} & {}^s J_{F,a_1}^{\frac{\alpha}{k}} h(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (g\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} g(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi\psi)(\tau) \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (g\psi)(\tau) \\ & \quad + {}^s J_{F,a_1}^{\frac{\beta}{k}} (g\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\psi)(\tau). \end{aligned} \tag{2.8}$$

Proof. Multiplying both sides of (2.4) by $\frac{F(v,s)g(v)}{k\Gamma_k(\beta)[\mathbb{F}(\tau,v)]^{1-\frac{\beta}{k}}}$, then we integrate the resulting inequality with respect to v over (a_1, τ) , we conclude that

$$\begin{aligned} & {}^s J_{F,a_1}^{\frac{\alpha}{k}} h(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (g\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} g(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi\psi)(\tau) \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (g\psi)(\tau) \\ & \quad + {}^s J_{F,a_1}^{\frac{\beta}{k}} (g\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (h\psi)(\tau). \end{aligned}$$

Thus, the Lemma is proved. □

Remark 2.6. If we take the kernel $F(t, s) = 1$ and $k = 1$ we obtain Lemma 6 of [4], and if $F(t, s) = h'(t)$, this result reduces to Lemma 2 of [9].

Theorem 2.7. Let φ, ψ be two synchronous functions on $[0, \infty)$ and let $f, \phi, \omega \geq 0$, then for all $\tau > a_1 \geq 0$, $\alpha, \beta > 0$, and $s \neq -1$, we get

$$\begin{aligned}
 & {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (f\varphi\psi)(\tau) + 2 {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi\psi)(\tau) \right. \\
 & \quad \left. + {}^s J_{F,a_1}^{\frac{\beta}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi\psi)(\tau) \right] \\
 & + \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} \phi(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \right] {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi\psi)(\tau) \\
 & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\psi)(\tau) \right] \\
 & + {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right] \\
 & + {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (f\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right]. \quad (2.9)
 \end{aligned}$$

Proof. Substituting in (2.8): $h = f$ and $g = \phi$, then we multiple the resulting inequality by ${}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau)$, we have

$$\begin{aligned}
 & {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi\psi)(\tau) \right] \\
 & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (f\psi)(\tau) \right]. \quad (2.10)
 \end{aligned}$$

Replacing again in (2.8): $h = \omega$ and $g = \phi$, and multiplying the result by ${}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau)$, we get

$$\begin{aligned}
 & {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} \phi(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi\psi)(\tau) \right] \\
 & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} f(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} (\phi\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right]. \quad (2.11)
 \end{aligned}$$

Now, by using the same idea, we put $h = \omega$ and $g = f$, and then we multiply the result by ${}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau)$, we conclude that

$$\begin{aligned}
 & {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} \omega(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (f\varphi\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} f(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi\psi)(\tau) \right] \\
 & \geq {}^s J_{F,a_1}^{\frac{\alpha}{k}} \phi(\tau) \left[{}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\beta}{k}} (f\psi)(\tau) + {}^s J_{F,a_1}^{\frac{\beta}{k}} (f\varphi)(\tau) {}^s J_{F,a_1}^{\frac{\alpha}{k}} (\omega\psi)(\tau) \right]. \quad (2.12)
 \end{aligned}$$

Finally, by adding the inequalities (2.10), (2.11) and (2.12) we obtain (2.9). □

Remark 2.8. In the case of the classical Riemann-Liouville Integral, this result extends the Theorem 4 of [4], and if we use the $(k; h)$ -Riemann-Liouville integral fractional, we obtain the Theorem 4 of [9].

Remark 2.9. Remark 2 and Remark 3 of [9], and Remark 7 of [4] are true for the general kernel used in our work.

3. Conclusions

In our work we obtained several generalized integral inequalities, which contain, as a particular case, some of those known in the literature, for example, if in Theorem 2.7 we consider $\alpha = \beta = 1$, $k = 1$ and $F(t, s) = 1$, we obtain the well-known Chebyshev Inequality (see [3]). In the same direction, we can add that one of the strengths of our results lies in the fact that by suitably choosing of F , i.e., if we consider kernels different than those indicated in the various remarks, one can further easily obtain additional integral inequalities involving the various types of fractional integral operators from our main results.

Acknowledgments. We would like to thank the referees for their careful reading of the manuscript and for some helpful suggestions that have improved the paper.

References

- [1] Abdeljawad, T., Mohammed, P.O., Kashuri, A., *New modified conformable fractional integral inequalities of Hermite-Hadamard type with applications*, J. Funct. Space., (2020), Article ID 4352357, 14 pages.
- [2] Baleanu, D., Mohammed, P.O., Vivas-Cortez, M., et al., *Some modifications in conformable fractional integral inequalities*, Adv. Differ. Equ., **2020**(2020), 374.
- [3] Chebyshev, P.L., *Sur les expressions approximatives des integrales dfinies par les autres prises entre les memes limite*, Proc. Math. Soc. Kharkov, **2**(1882), 93-98.
- [4] Dahmani, Z., *New Inequalities in Fractional Integrals*, International Journal of Nonlinear Science, **9**(2010), 493-497.
- [5] Díaz, R., Pariguan, E., *On hypergeometric functions and Pochhammer k-symbol*, Divulg. Mat., **15**(2007), no. 2, 179-192.
- [6] Fernandez, A., Mohammed, P., *Hermite-Hadamard inequalities in fractional calculus defined using Mittag-Leffler kernels*, Math. Meth. Appl. Sci., (2020), 1-18, <https://doi.org/10.1002/mma.6188>.
- [7] Galeano, J.D., Nápoles, J.E., Pérez, E., *On a general formulation of the fractional operator Riemann-Liouville and related inequalities*, (submitted).
- [8] Guzman, P.M., Karus, P., Nápoles Valdes, J.E., *Generalized integral inequalities of Chebyshev type*, Fractal Fract., (2020), no. 4, 10, doi:10.3390/fractalfract4020010.
- [9] Mohammed, P.O., *Inequalities of $(k; s)$, $(k; h)$ -type for Riemann-Liouville fractional integrals*, Applied Mathematics E-Notes, **17**(2017), 199-206.
- [10] Mohammed, P.O., Abdeljawad, T., *Opial integral inequalities for generalized fractional operators with nonsingular kernel*, J. Inequal. Appl., **2020**(2020), 148.
- [11] Mohammed, P.O., Abdeljawad, T., *Integral inequalities for a fractional operator of a function with respect to another function with nonsingular kernel*, Adv. Differ. Equ., **2020**(2020), 363.

- [12] Mohammed, P.O., Brevik, I., *A new version of the Hermite-Hadamard inequality for Riemann-Liouville fractional integrals*, Symmetry, **12**(2020), 610, doi:10.3390/sym12040610.
- [13] Mohammed, P.O., Sarikaya, M.Z., *On generalized fractional integral inequalities for twice differentiable convex functions*, J. Comput. Appl. Math., **372**(2020), 112740.
- [14] Mubeen, S., Habibullah, G.M., *k-fractional integrals and applications*, Int. J. Contemp. Math. Sci., **7**(2012), 89-94.
- [15] Mubeen, S., Habibullah, G.M., Naeem, M.N., *The Minkowski inequality involving generalized k-fractional conformable integral*, J. Inequal. Appl., **81**(2019), <https://doi.org/10.1186/s13660-019-2040-8>.
- [16] Nápoles Valdes, J.E., Rodriguez, J.M., Sigarreta, J.M., *New Hermite-Hadamard type inequalities involving non-conformable integral operators*, Symmetry, (2019), 11, 1108, doi:10.3390/sym11091108.
- [17] Qi, F., Guo, B.N., *Integral representations and complete monotonicity of remainders of the Binet and Stirling formulas for the gamma function*, Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat., **111**(2017), no. 2, 425-434, <https://doi.org/10.1007/s13398-016-0302-6>.
- [18] Qi, F., Habib, S., Mubeen, S., Naeem, M.N., *Generalized k-fractional conformable integrals and related inequalities*, AIMS Mathematics, **4**(3)(2019), 343-358, DOI:10.3934/math.2019.3.343.
- [19] Rainville, E.D., *Special Functions*, Macmillan Co., New York, 1960.
- [20] Rashid, S., Hammouch, Z., Kalsoom, H., Ashraf, R., Chu, Y.M., *New investigation on the generalized k-fractional integral operators*, Frontiers in Physics, **8**(2020), Article 25, doi: 10.3389/fphy.2020.00025.
- [21] Sarikaya, M.Z., Dahmani, Z., Kiris, M.E., Ahmad, F., *(k, s)-Riemann-Liouville fractional integral and applications*, Hacettepe Journal of Mathematics and Statistics, **45**(2016), no. 1, 77-89.
- [22] Yang, Z.H., Tian, J.F., *Monotonicity and inequalities for the gamma function*, J. Inequal. Appl., **317**(2017), <https://doi.org/10.1186/s13660-017-1591-9>.
- [23] Yang, Z.H., Tian, J.F., *Monotonicity and sharp inequalities related to gamma function*, J. Math. Inequal., **12**(2018), no. 1, 1-22, <https://doi.org/10.7153/jmi-2018-12-01>.

Juan Gabriel Galeano Delgado

Facultad de Ciencias e Ingeniería, Universidad del Sinú Elias Bechara Zainúm,
Monteria, Colombia

e-mail: juangaleanod@unisinu.edu.co

Juan E. Nápoles Valdés

UNNE, FaCENA, Ave. Libertad 5450, Corrientes 3400, Argentina

e-mail: jnapoles@exa.unne.edu.ar

UTN-FRRE, French 414, Resistencia, Chaco 3500, Argentina

e-mail: jnapoles@frre.utn.edu.ar

Edgardo Pérez Reyes

Facultad de Ciencias e Ingeniería, Universidad del Sinú Elias Bechara Zainúm,
Monteria, Colombia

e-mail: edgardomath@gmail.com

